

## Non-extractable polyphenols, a major dietary antioxidant: occurrence, metabolic fate and health effects

Jara Pérez-Jiménez, M. Elena Díaz-Rubio and Fulgencio Saura-Calixto\*

*Department of Metabolism and Nutrition, Institute of Food Science, Technology and Nutrition (ICTAN-CSIC), c/ José Antonio Novais, 10, 28040 Madrid, Spain*

### Abstract

Current research on dietary antioxidants misses the so-called non-extractable polyphenols (NEPP), which are not significantly released from the food matrix either by mastication, acid pH in the stomach or action of digestive enzymes, reaching the colon nearly intact. NEPP, not detected by the usual analytical procedures, are made up of macromolecules and single phenolic compounds associated with macromolecules. Therefore, NEPP are not included in food and dietary intake data nor in bioavailability, intervention or observational studies. The present paper aims to provide an overview of dietary NEPP – nature, occurrence in diet, metabolic fate and possible health effects. NEPP are a relevant fraction of dietary polyphenols exerting their main biological action in the colon, where they are extensively fermented by the action of microbiota, giving place to absorbable metabolites. NEPP exhibit different potential health-related properties, in particular in relation to gastrointestinal health, such as increases in antioxidant and antiproliferative capacities, reduction of intestinal tumorigenesis and modification of gene expression, as observed in different animal models. Further research into NEPP may provide a better understanding of the health effects of dietary antioxidants.

**Key words:** Polyphenols: Non-extractable polyphenols: Dietary antioxidants: Dietary intake: Gut health

### Introduction

There is increasing scientific evidence of the existence of a relationship between diets rich in natural antioxidants and the prevention of several chronic diseases. Epidemiological studies have shown inverse associations between total dietary antioxidant intake and inflammation<sup>(1,2)</sup>, ischaemic stroke<sup>(3)</sup>, alterations in endothelial function<sup>(4)</sup> and gastric cancer<sup>(5)</sup>. Moreover, adherence to a Mediterranean diet, which implies a higher plasma concentration of dietary antioxidants<sup>(6)</sup>, is associated with a decrease in overall mortality and particularly in mortality resulting from CVD or cancer<sup>(7)</sup>.

Dietary antioxidants are a diverse group of chemical compounds, with varying solubility in biological fluids or biomembranes, and mechanisms of actions; they include carotenoids, vitamins C and E, and polyphenols<sup>(8)</sup>. Of these, polyphenols (which include a wide variety of chemical structures that share one or more phenol groups) are the most commonly consumed dietary antioxidants, corresponding to about 90% of dietary antioxidant intake<sup>(9,10)</sup>.

The interest in polyphenol research within the fields of nutrition and food science has led to an increasing literature dealing with: (a) the bioavailability and distribution

in tissues of polyphenols<sup>(11)</sup>; (b) possible mechanisms of action of polyphenols<sup>(12)</sup>; and (c) the health effects derived from polyphenol intake<sup>(13)</sup>. In particular, clinical trials and/or epidemiological studies have shown that polyphenols play a role in the prevention of CVD and of certain kinds of cancer<sup>(14,15)</sup>.

Current research into polyphenols focuses mainly on only a fraction of dietary polyphenols, corresponding to those that can be extracted from food with aqueous–organic solvents. The polyphenols identified in such extracts – named extractable polyphenols (EPP) – are then considered to be the total polyphenol content, and are used as the basis for calculations of dietary intake, bioavailability studies and to design intervention or observation studies.

However, it is known that a significant fraction of food polyphenols remains in the corresponding residues after the extraction; the so-called non-extractable polyphenols (NEPP)<sup>(6)</sup>. Since foodstuffs are consumed as a whole, i.e. including both EPP and NEPP, significant amounts of NEPP are ingested daily, contributing to the reported health effects of polyphenols. However, this fraction of dietary polyphenols has long been neglected and studies of it are still scarce. The present review aims to provide

**Abbreviations:** DF, dietary fibre; EPP, extractable polyphenols; NEPA, non-extractable proanthocyanidins; NEPP, non-extractable polyphenols.

\*Corresponding author: Dr Fulgencio Saura-Calixto, fax +34 91 549 36 27, email [fsaura@ictan.csic.es](mailto:fsaura@ictan.csic.es)

an overview of dietary NEPP: their nature, occurrence in the diet, metabolic fate and possible health effects.

### Non-extractable polyphenols

Several classifications for the different polyphenol classes, focused exclusively on EPP, have been proposed based on the different chemical skeletons that they present. One of the most common classifications divides them into flavonoids, phenolic acids, stilbenes, lignans and other polyphenols (including tyrosols and alkylresorcinols). All of them possess at least one aromatic ring, with one or more hydroxyl moieties. Several procedures have been reported for extracting polyphenols from plant foods; the most common is solid–liquid extraction with different combinations of organic solvents (methanol, ethanol or acetone) with water. Since there is abundant literature on EPP (more than 30 000 references according to the Thomson Reuters (formerly ISI) Web of Knowledge website; <http://wokinfo.com/>), including aspects such as their chemical characterisation, bioavailability, nutritional qualities and effects on health, this class of polyphenols is beyond the scope of the present review.

When performing such chemical extractions on foodstuffs, a solid residue remains. In polyphenol research, the residue is usually ignored and considered to be devoid of polyphenols. However, the residue is actually a rich source of another fraction of polyphenols with specific biological activities: NEPP.

NEPP are those dietary polyphenols that, after ingestion, are not significantly released from the food matrix either by mastication, an acidic pH in the stomach or the action of digestive enzymes. They reach the colon nearly intact, where they are subjected to extensive transformation by colonic microflora (see ‘Non-extractable polyphenols through the human gut’ section). NEPP include macromolecules, such as high-molecular-weight proanthocyanidins, and single phenolic compounds, such as phenolic acids, associated with macromolecules, mainly polysaccharide constituents of dietary fibre (DF) and protein. They are generally not included in polyphenol analysis of foodstuffs, although their NEPP content may be even higher than that of the EPP fraction (see ‘Occurrence of non-extractable polyphenols in foodstuffs and diets’ section). With regards to their chemical nature, NEPP comprise mainly polyphenols also found as EPP, such as proanthocyanidins, other flavonoids, phenolic acids and hydrolysable tannins<sup>(16)</sup>.

NEPP interact with the food matrix (mainly polysaccharides and proteins) via different mechanisms: (a) hydrogen bonding, as described for non-extractable proanthocyanidins (NEPA)<sup>(17)</sup> and that may also be applicable to hydrolysable tannins; (b) hydrophobic interactions, including possible encapsulation into hydrophobic pockets with NEPA<sup>(17)</sup>; (c) covalent bonding (to form esters and ether bonds), in the case of phenolic acids<sup>(18–20)</sup>, and possibly of NEPA and hydrolysable tannins. In addition, a fraction

of EPP may be associated with NEPP<sup>(21)</sup>. Whatever the case regarding that last point, it should be pointed out that there are still many aspects related to the interactions, as well as to the physico-chemical structures of NEPP, that have yet to be elucidated.


Another important aspect of NEPP is that they are associated with DF and they can be considered to be constituents of DF. Since the first research at the end of the 1980s, which showed that this fraction of dietary polyphenols were constituents of DF<sup>(22,23)</sup>, in recent years some papers have emphasised the significant contribution of NEPP to the health-related properties attributed to DF polysaccharides<sup>(24–26)</sup>. Therefore, NEPP do indeed contribute to some of the health-related properties commonly attributed to DF and EPP.

### Occurrence of non-extractable polyphenols in foodstuffs and diets

Figure 1 shows the process followed for the analysis of the NEPP content of food<sup>(27–29)</sup>. A chemical extraction with aqueous–organic solvents is carried out on the food, which releases EPP in the supernatant fraction and also produces a residue. The residue is then subjected to the action of digestive enzymes, to obtain a new residue containing NEPP. Chemical hydrolysis is then carried out, in order to release NEPP from the food matrix, or to fragment the polymers. The hydrolysates obtained after these treatments may then be analysed using spectrophotometric or chromatographic techniques. Although the NEPP content of food has been studied much less than the EPP content, the available data show the relevance of this fraction of dietary polyphenols.

### Occurrence of non-extractable polyphenols in individual foodstuffs

The different classes of NEPP are widespread among the different families of plant foods (cereals, fruit, vegetables,



**Fig. 1.** Scheme of the analysis of food non-extractable polyphenols (NEPP). EPP, extractable polyphenols; DAD, diode array detection.

nuts, legumes), although some of them are particularly characteristic of a particular food family. Table 1<sup>(27,29–53)</sup> shows the reported content of NEPP belonging to different classes (proanthocyanidins, hydrolysable tannins and hydrolysable phenolics) in several plant foods.

NEPA are mostly found in fruit. Despite being obtained via different procedures, the data included in Table 1 provide a rough estimation of the NEPA content of foodstuffs. The case of the banana is particularly interesting, since its NEPA content (recently confirmed by an improved procedure<sup>(54)</sup>) would be about 100-fold the reported extractable proanthocyanidin content for this food<sup>(55)</sup>. This reflects the contribution of this fraction of dietary polyphenols, that is usually ignored, to the total content of polyphenolic compounds in foodstuffs and therefore to dietary intake of such compounds.

Hydrolysable phenolics are comprised of several classes of phenolic compounds, although the most common are phenolic acids (ferulic acid, caffeic acid, sinapic acid and others). Although hydrolysable phenolics have usually only been studied in cereal products, they have recently

been reported to be present in many other foodstuffs, such as onion, black olive, apple, medlar, mandarin, acerola, cashew apple, black currant pomace and red ginseng, as shown in Table 1.

Finally, hydrolysable tannins are more specific to certain foodstuffs. Ellagic acid and other derivatives from the hydrolysis of ellagitannins have been reported, for instance, to be present in several nuts (Table 1).

Another problem when it comes to evaluating the contribution of NEPP to the total polyphenol content of foodstuffs is that besides the fact that studies dealing with the NEPP content of foodstuffs are rare, they usually focus on only one class of NEPP (as is the case with most of the references included in Table 1) and do not consider the overall analysis of the different classes of NEPP. However, some methods have been proposed that include both the analysis of hydrolysable phenolics and NEPA in the evaluation of NEPP<sup>(29,56)</sup>; some of the results are shown in Fig. 2<sup>(57)</sup>. The analysis of EPP, hydrolysable phenolics and NEPA in several types of fruit and nuts showed that the contribution of NEPP was between 60 and 90 % of the total polyphenol content, and therefore represented the major fraction of these dietary antioxidants.

Finally, since NEPP are constituents of DF, updated methods for the analysis of DF include analysis of NEPP<sup>(37)</sup>. Such an approach has already been used in the analysis of DF from several types of fruit, vegetables and nuts<sup>(30,37,41)</sup>.

### Occurrence of non-extractable polyphenols in diets

Analysis of NEPP in individual foodstuffs should serve as the basis for calculating NEPP content in whole diets in different populations; and therefore, for estimating NEPP intake from them. A few papers have dealt with the analysis and dietary intake of NEPP in the Spanish diet, as an example of the Mediterranean diet<sup>(28,56)</sup> and in a rural Mexican population<sup>(58)</sup>.

As regards the Spanish diet<sup>(28,56)</sup> when determining total polyphenol content in the different food consumed, it was found that in all the plant-based food groups (cereals, vegetables, legumes, fruit and nuts) the NEPP content was higher than that of EPP. When the data were translated into estimates of intake, total polyphenol intake in the Spanish diet was estimated to be about 1800–3000 mg/individual per d, depending on the analytical techniques used, with NEPP contributing about 50 % of the total polyphenol intake. Fruits were the highest contributor to NEPP intake (about 47 %), followed by cereals (31 %), vegetables (13 %), legumes (6 %) and nuts (4 %).

As the authors of the studies emphasised, some of the data are closer to estimates than to accurate determinations of NEPP intake, due to the analytical limitations. Nevertheless, the results constitute evidence of an appreciable presence of NEPP in plant foods and of their contribution to total polyphenol intake.


**Table 1.** Non-extractable polyphenol content reported in the literature for several plant foods

Non-extractable polyphenol class	Food	Content (mg/100 g fw)	Reference
Proanthocyanidins	Açaí fruit	1240 (SD 140)*	(30)
	Apple	37–43	(31)
	Quince	48	(32)
	Banana	980 (SD 45)*	(27)
	Apple pomace	18–23	(33)
	Cranberry pomace	1685	(34)
	Cocoa powder	602 (SD 13)	(35)
	Carob pod	180	(36)
	Onion	410 (SD 20)*	(37)
	Black olive	14–40	(38)
Hydrolysable phenolics	Apple	78 (SD 6)	(29)
	Medlar	0.5–1.0	(39)
	Mandarin	39–107	(40)
	Acerola	390 (SD 10)*	(41)
	Cashew apple	1210 (SD 70)*†	(41)
	Black currant pomace	41 (SD 1)	(42)
	Red ginseng	2.0 (SD 0.1)	(43)
	Refined maize flour	209	(44)
	Whole-grain maize flour	174 (SD 2)	(45)
	Whole-grain wheat flour	32	(46)
	Wheat bran	26	(47)
	Rice bran	243*	(48)
	Whole-grain barley flour	60–135	(49)
	Triticale bran	137 (SD 1)	(50)
	Millet	32–168	(51)
Hydrolysable tannins	Brazil nut	210	(52)
	Heartnut	115	(53)

fw, Fresh weight.

\*Results expressed per 100 g dry weight.

†Spectrophotometric analysis, so may include gallic acid and/or ellagic acid derived from hydrolysable tannins.



**Fig. 2.** Contribution of the different classes of non-extractable polyphenols to total polyphenol content in different foods<sup>(29,41,57)</sup>. ■, Non-extractable proanthocyanidins; ■, hydrolysable phenolics; ■, extractable polyphenols.

Another study focused specifically on the contribution of NEPP to polyphenol intake from fruit and vegetables in a rural Mexican population<sup>(58)</sup>. Despite the food items included in the categories in Mexico being very different from those consumed in Spain, it is remarkable that the results again showed that NEPP are more abundant in the solid plant-based food groups than EPP are and, therefore, they contribute significantly to the total polyphenol intake.

This kind of evaluation of total polyphenol content in diets, including NEPP, as well as estimations of the associated polyphenol intake, should be generalised to different populations as a starting point to advance our knowledge of the possible associations between NEPP intake and health effects that have not previously been considered.

### Non-extractable polyphenols through the human gut

To understand the possible health effects associated with the intake of any bioactive compound, and therefore of NEPP, it is necessary to have knowledge of the different events that occur along the human gut, including:

- (1) The solubilisation of the compounds in intestinal fluids, via different mechanisms, resulting in the compounds becoming bioaccessible. This may be estimated by determining the presence of NEPP in intestinal fluids, both in the small and in the large intestine.
- (2) Possible transformations by colonic microbiota of the compounds that are bioaccessible in the colon. Such transformations may be enhanced or inhibited by the presence of other food components and/or other interactions with the microbiota. The metabolites


derived from such transformations may be determined either *in vitro* (supernatant fractions from *in vitro* fermentation) or *in vivo* (caecal contents or faeces).

- (3) Absorption, either of the original compounds or of the derived metabolites. Absorption gives rise to metabolites that may be determined in different biological fluids, such as urine and blood.
- (4) Possible effects on target tissues. An accumulation of bioactive metabolites in certain tissues may contribute to the local effects of compounds of interest.

All these aspects have been studied in depth over recent years for EPP<sup>(11)</sup>, but studies on NEPP are scarce. Despite that, the available work that focuses on NEPP suggests that they are subjected to extensive transformation in the gut, giving rise to several absorbable metabolites that have been reported to have different health effects. From such work<sup>(28,59–65)</sup> a general scheme of the metabolic fate of NEPP can be proposed, as shown in Fig. 3, where compound 'a' represents a hydrolysable phenolic compound, i.e. a phenolic acid associated with a polysaccharide chain, and compound 'b' represents a polymeric NEPA. Briefly, most NEPP will pass through the small intestine (Fig. 3(A)) without any transformation. Once in the colon (Fig. 3(B)), and mainly through the action of the microbiota, NEPP are released and new compounds form. Some of these microbial metabolites may be absorbed through the portal vein, reaching the liver (Fig. 3(C)), where several related processes occur, giving rise to phase II metabolites. Once formed, these metabolites may return to the digestive tube through the bile (Fig. 3(D)), or pass into the bloodstream as a first step towards reaching the target tissues and finally be excreted in urine (Fig. 3(E)).

### Bioaccessibility of non-extractable polyphenols

Bioaccessibility corresponds to the amount of a food constituent that is solubilised in intestinal fluids as a consequence of physico-chemical conditions, or the action of either digestive enzymes or bacterial microbiota. Saura-Calixto *et al.*<sup>(28)</sup> evaluated the bioaccessibility of total polyphenols, including NEPP, in a complete diet by using an *in vitro* gastrointestinal model followed by *in vitro* colonic fermentation. They observed that, while about a 50% of hydrolysable phenolics become bioaccessible in the small intestine, in the case of NEPA most of them arrive nearly intact to the colon. Similarly, other studies have shown that NEPA are only partially depolymerised in the small intestine<sup>(64,65)</sup>. Once in the colon, those NEPP that are not solubilised in the small intestine become accessible either by: (a) fermentation by colonic microbiota of the molecules to which they are associated (carbohydrates or proteins); or (b) the action of some intestinal enzymes able to break covalent bonds, such as esterases<sup>(59)</sup>. It has been estimated that 95% of NEPA arriving to the colon are released from the food matrix by these two mechanisms<sup>(28)</sup>.



**Fig. 3.** Overview of the metabolic fate of non-extractable polyphenols (NEPP). Sulf-, sulfur; Me-, methyl; Gluc-, glucose.

Therefore, significant amounts of the different classes of NEPP appear in the large intestine daily as bioaccessible compounds, including both the compounds that have been solubilised in the small intestine but have not been absorbed, as well as those released in the large intestine. Since NEPP are widespread among the different groups of plant foods, a diverse diet including food from all of them would guarantee a continuous supply of beneficial bioaccessible compounds through the digestive tract. Obviously, a fraction of EPP not previously absorbed in the small intestine also arrives as bioaccessible compounds to the large intestine<sup>(28)</sup>; future work should elucidate the particular contributions of EPP and NEPP to the pool of bioaccessible phenolic compounds present in the large intestine.

### Colonic transformation and bioavailability of non-extractable polyphenols

In the colon, NEPP released from the food matrix and polymeric NEPP undergo colonic fermentation, which produces potentially absorbable metabolites. Work focusing strictly on the fermentation of dietary NEPP is still scarce and mostly qualitative<sup>(62–65)</sup>, but it has already

provided clear evidence on their colonic fermentation. For instance, supplementing rats with a NEPP concentrate devoid of EPP resulted in the presence of polyphenol-derived metabolites in the urine and faeces (ten and three, respectively) at concentrations at least 10-fold higher than in a non-supplemented group<sup>(65)</sup>.

On the other hand, since many studies of the metabolic fate of polyphenols are based on supplementation with whole foods, many of which evidently contain NEPP, some of the data regarding the metabolites derived from polyphenols would actually correspond to NEPP metabolites. Therefore, considering specific studies on NEPP fermentation<sup>(62–66)</sup> and other studies on foods containing both EPP and NEPP<sup>(67–69)</sup>, these would be the main microbial metabolites derived from the different fractions of NEPP<sup>(62–69)</sup>:

- (1) Metabolites derived from NEPA. These include: (a) the monomeric constituents of NEPA polymers, mainly (epi)catechin; and (b) a wide variety of compounds derived from the breakdown of the basic skeleton of flavanols and further degradation, which gives place first to valerolactones, then to several phenolic acids corresponding to successive degradations





- (hydroxyphenylpropionic acids, hydroxyphenylacetic acids, etc.) and then to small acids such as hippuric acid.
- (2) Metabolites derived from hydrolysable phenolics. Although free phenolic acids released from the food matrix may be absorbed directly, they may also undergo additional transformation resulting from interaction with the colonic microbiota. For instance, ferulic acid is transformed into dihydroferulic acid, which is later metabolised to other phenolic acids coincident with the degradation of proanthocyanidins.
  - (3) Metabolites derived from hydrolysable tannins. Ellagic acid released from ellagitannins in the gastrointestinal tract is transformed by the microbiota into a particular class of compounds, the urolithins.

The presence of these compounds has been determined in caecal contents and in faeces<sup>(64,65)</sup>, but their determination is most common in urine and blood (in both animal studies and clinical trials)<sup>(62,63,66)</sup>. The results show that these compounds are absorbed and therefore become bioavailable and can have different biological effects on the human body, as well as possible local effects on the colon (see 'Health-related properties of non-extractable polyphenols' section).

The metabolites described above do not differ from those described as derived from EPP colonic fermentation<sup>(11,68)</sup>, i.e. the colonic fermentation of both EPP and NEPP gives place to the same compounds. However, the specific characteristics of NEPP (high molecular weight and/or association with other macromolecules present in the food matrix) cause two specific features in the colonic transformation of NEPP as compared with EPP:

- (1) The absorption of NEPP is retarded in relation to that of EPP, as shown by the delayed peak in plasma ferulic acid levels after the intake of wheat bran (containing NEPP) in comparison with that of pure ferulic acid<sup>(60)</sup>, or the delayed peak in plasma antioxidant capacity after the intake of a NEPP-rich matrix in comparison with the data for EPA-rich matrixes<sup>(40)</sup>. Such data indicate that metabolites derived from NEPP circulate for longer periods in the human body than those derived from EPP. Indeed, some of the metabolites detected in the urine and faeces of rats fed with NEPP-rich products and in the supernatant fractions from *in vitro* fermentation of these matrixes were also detected in plasma from fasting volunteers after 24 h of a polyphenol-free diet, which demonstrates the prolonged circulation times of NEPP metabolites<sup>(63)</sup>. Therefore, NEPP may continually provide the colonic microbiota with a significant amount of fermentable substrates that need more time to be released and fermented than EPP due to their specific nature, and, therefore, result in a more prolonged circulation of beneficial metabolites.
- (2) Since NEPP do not reach the colon alone, but in combination with other fermentable dietary components,

such as proteins and polysaccharides, their fermentation rate may be affected. Indeed, some studies in rats have shown that the presence of DF enhances the fermentation of polyphenols<sup>(70,71)</sup>. Moreover, in the work by Saura-Calixto *et al.*<sup>(63)</sup>, the *in vitro* fermentability (as SCFA production compared with a control) of a concentrate of polymeric NEPP was 23%, while that of a concentrate of NEPP associated with DF was 50%, probably due to an increase in bacterial activity. Therefore, the colonic fermentation of NEPP would be accompanied by a synergism with the other components of the food matrix to which they are associated, an aspect that is not present in the fermentation of EPP, which are free in the food matrix.


A summary of some of the colonic metabolites derived from NEPP, as well as the specific features of their fermentation, is shown in Fig. 4. Nevertheless, more specific work on the colonic metabolism of NEPP from different origin and/or different nature, including quantitative data, is needed. This would allow elucidation of the relative contributions of EPP and NEPP to the dozens of polyphenol-derived metabolites that circulate in the human body after plant food intake, and that may be indeed be responsible for many of the reported benefits of polyphenols<sup>(69)</sup>.

### Health-related properties of non-extractable polyphenols

As previously described, NEPP undergo extensive transformation in the colon. Therefore, the health effects associated with them may not come mainly from the intact NEPP, but from their metabolites. In recent years, a number of studies have started to focus on the biological activities of polyphenol metabolites. Although they do not specifically consider the contribution of NEPP to their release, a significant proportion of the production of such compounds in the human body, and therefore of their associated effects, would be due to NEPP intake, as explained above.

Table 2<sup>(72–76)</sup> summarises some of the findings regarding the biological effects of these metabolites; the results are mostly obtained in cell cultures and animal models. The effects include anti-inflammatory activity, a reduction of oxidative stress, the inhibition of protein glycation, antiproliferative effects and effects on flow-mediated vasodilation<sup>(72–76)</sup>. These are probably the mechanisms that underlie the observed health effect of NEPP-rich products in animal and human studies.

Moreover, since a significant fraction of NEPP is present in foods associated with polysaccharides, when such a complex reaches the colon, the polysaccharide fraction also ferments, releasing beneficial components such as butyric acid, associated with the prevention of colorectal cancer<sup>(77,78)</sup>, and acetate and propionate, with potential health effects on lipid metabolism<sup>(78,79)</sup>. These metabolites also contribute to the health effects reported in animal or human studies of NEPP-rich products.



**Fig. 4.** Main features of non-extractable polyphenol (NEPP) colonic fermentation. EPP, extractable polyphenols.

### Gastrointestinal health

Since NEPP reach the colon nearly intact and that is where they suffer most of their metabolic transformation, it is evident that one of their major targets of action is gastrointestinal health. Several studies have shown promising results in this field after supplementation of different animal models with several NEPP-rich products (Table 3)<sup>(80–91)</sup>. Grape antioxidant DF, a matrix derived from wine production with a high content of NEPA<sup>(92)</sup>, mostly associated with DF, has been the most commonly tested matrix; however, some studies have also fed rats with apple pulp, carob pod, artichoke, grape pomace or

grape seeds. Most studies included ten animals per group (commonly Wistar rats) and carried out the supplementation by including a 5–13% of the tested product in the diet. The duration of the study was commonly 4 weeks.

Overall, these animal studies show that NEPP may have a beneficial effect on gastrointestinal health through a combination of mechanisms<sup>(80–91)</sup>:

- (1) An increase in stool weight, reducing transit time and therefore the time of contact of toxic compounds with the colonic epithelium<sup>(80–83)</sup>.
- (2) An increase in the antioxidant capacity of caecal content and in the expression of endogenous antioxidant

**Table 2.** Summary of biological effects described for polyphenol metabolites

Metabolites	Biological effect	Reference
(–)-Epicatechin conjugates	Effects on flow-mediated vasodilation in human subjects	(72)
3-Phenylpropionic acid, 3-hydroxyphenylacetic acid, 3-hydroxyphenylpropionic acid	COX-2 inhibition in HT-29 intestinal cells	(73)
3,4-Dihydroxyphenylacetic acid, hydroferulic acid	Anti-inflammatory activity in CDD-18 colon fibroblast cells	(74)
Hydrocaffeic acid	Reduction of oxidative stress in DSS-treated rats	(74)
Urolithin A, urolithin B, dihydroferulic acid	Protein glycation inhibition	(75)
Urolithin A, urolithin B	Antiproliferative effects through change in the expression levels of growth factor receptors, oncogenes and tumour suppressors in Caco-2 intestinal cells	(76)

COX-2, cyclo-oxygenase-2; DSS, dextran sodium sulfate.



**Table 3.** Summary of animal studies on the effect of non-extractable polyphenols (NEPP) on gastrointestinal health

NEPP-rich product	Daily dose and duration of the study	Animal model and size of treatment group	Reported effects	Reference
Apple pulp, carob pod, grape pomace, grape seeds	10 % of the diet, 1–8 weeks	Wistar rats, <i>n</i> 10–12	Increase in stool weight (+50 to +300 %)	(80–83)
Artichoke, grape seeds	5–13 % of the diet, 3–5 weeks	Wistar rats, <i>n</i> 10	Increase in antioxidant capacity in caecal content (+400 to +1000 %)	(84,85)
Grape antioxidant dietary fibre	5 % of the diet, 4 weeks	Wistar rats, <i>n</i> 10	Reduction of lipid oxidation in distal colonic mucosa (–25 %)	(86)
Grape antioxidant dietary fibre	0.1 g/kg bw, 2 weeks	C57BL/6J mice, <i>n</i> 8	Overexpression of enzymes pertaining to the xenobiotic detoxifying system and endogenous antioxidant cell defences (+100 to +200 %)	(87)
Grape antioxidant dietary fibre	0.1 g/kg bw, 2 weeks	C57BL/6J mice, <i>n</i> 8	Down-regulation in colon mucosa of genes associated with tumour development and proto-oncogenes (–100 to –200 %), up-regulation of tumour-suppressor genes (+100 %)	(87)
Grape antioxidant dietary fibre	1 % of the diet, 16 weeks	APC <sup>Min/+</sup> mice, <i>n</i> 12	Reduction in the number (–76 %) and size (–65 to –87 %) of polyps	(88)
Grape antioxidant dietary fibre	5–7 % of the diet, 4 weeks	Wistar rats, <i>n</i> 10	Induction of epithelial hypoplasia in colonic mucosa (–30 % DNA content), reduction of apoptosis (–30 to –40 %), decrease of number of crypts (–10 to –15 %)	(86,89,90)
Grape pomace, grape antioxidant dietary fibre	5 % of the diet, 4 weeks	Wistar rats, <i>n</i> 10	Stimulation of <i>Lactobacillus</i> growth (1-log increase)	(91)

bw, Body weight.

systems, thus counteracting the tumour-promoting reactive oxygen species (ROS) present in the colon<sup>(84,85,89,90)</sup>.

- (3) A prebiotic effect proven by stimulation of *Lactobacillus*, which would also imply a reduction in the proportion of non-beneficial bacteria species<sup>(91)</sup>. This prebiotic effect was observed when rats were supplemented with a NEPP-rich product but not when their diet contained the same amount of cellulose, indicating that this was a specific effect of NEPP and not of DF. Previous *in vitro* studies had also reported a prebiotic effect for grape pomace, another NEPP-rich product<sup>(92)</sup>.
- (4) An increase in antiproliferative capacity in healthy rats, shown by an induction of epithelial hypoplasia, reduction of apoptosis and a decrease in the number of crypts<sup>(86,89,90)</sup>. These antiproliferative effects were additionally found in cell cultures treated with a NEPP concentrate from apple pomace<sup>(70)</sup>.
- (5) A reduction of intestinal tumorigenesis in APC<sup>Min/+</sup> mice, an animal model of colorectal cancer, including significant reductions in the number (–76 %) and size (–65 to –87 %) of polyps<sup>(88)</sup>.
- (6) Modifications in gene expression in healthy mice and in APC<sup>Min/+</sup> mice, in particular, down-regulation of genes associated with tumour development and proto-oncogenes (for example, genes belonging to the *RAS* family, such as *RASSF4*, *RAP2C* and *RAP2B*), and up-regulation of tumour-suppressor genes (for instance, *NLB1* or *TGFb3*, related to cell cycle and cell proliferation, respectively)<sup>(87,88)</sup>.

In none of these studies neither adverse nor toxic effects were observed in the animals.

Overall, all these animal studies have shown a promising role for NEPP in the prevention and/or treatment of colorectal cancer and other gastrointestinal disorders, which should be validated in human subjects with proper clinical trials. Interestingly, when grape antioxidant DF was provided to human subjects, an increase in weekly faecal outputs was observed in those subjects who initially exhibited seven or fewer faecal outputs per week<sup>(93)</sup>. Although grape antioxidant DF is a DF-rich product and this bulking effect is well known for DF, it should be emphasised that NEPP are indeed constituents of DF<sup>(26)</sup>, although they are usually not considered, and they may make a specific contribution to this bulking effect.

Also, a recent epidemiological study<sup>(15)</sup> showed an inverse association between proanthocyanidin intake and the risk of colorectal cancer; furthermore the reduction in risk was greater, the higher the degree of polymerisation of the proanthocyanidins (> 10). This study only considered the intake of EPA, but since NEPA are also present in common foodstuffs and they consist of large polymeric structures, these results highlight the need to corroborate the possible role of NEPP – and in particular NEPA – in the prevention of colorectal cancer.

### CVD risk factors

The absorption of bioactive metabolites generated in the colon (both from the fermentation of NEPP themselves and from that of their associated polysaccharides) may



give rise to systemic effects that manifest in other organs and tissues. In particular, the possible role that these colonic metabolites may play in the prevention of CVD has been expressed as the 'gut–heart axis' hypothesis<sup>(94)</sup>. Several studies in animal models and/or human subjects<sup>(82,87,93,95)</sup> have shown that a NEPP-rich matrix may mitigate certain risk factors for CVD, such as hyperlipidaemia or hypertension. As regards to the effects in lipidaemia, this could occur through different mechanisms, such as a reduction in lipid biosynthesis (proven by a down-regulation of the expression of genes involved in this process) and an increase in faecal excretion of cholesterol<sup>(82,83,87)</sup>. Moreover, NEPP could also be related to the prevention of CVD by a reduction of lipid oxidation<sup>(93)</sup>, which has shown to be associated with CVD<sup>(96)</sup>.

Also, it should be considered that the association of NEPP with DF may give rise to specific synergies. In this way, the daily supplementation to hypercholesterolaemic subjects with 7.5 g of a grape-derived NEPP-rich product for 16 weeks resulted in significant reduction in plasma cholesterol and blood pressure, which were greater than those described separately for DF and EPP in several meta-analyses. This may be due to the combined effects of a single matrix containing DF and associated NEPP<sup>(92)</sup>.

Finally, a field which has so far not been explored and in which NEPP may have a preventive effect is that of the metabolic syndrome and, in particular, in the maintenance of homeostatic glucose levels through action on insulin sensitivity. The existence of metabolic crosstalk between the colon and periphery organs affecting insulin sensitivity has been suggested<sup>(97)</sup>. More recently, the supplementation of mice with grape antioxidant DF produced a decrease in blood glucose, compared with controls, and up-regulation of the gene encoding the enzyme glucose-6-phosphatase, which plays a key role in the homeostatic regulation of blood glucose<sup>(87)</sup>.

### Concluding remarks

NEPP constitute an important fraction of dietary polyphenols, which has commonly been ignored in polyphenol research.

The different classes of NEPP are widespread among all the families of plant foods, and their content in many foodstuffs is higher than that of EPP. Therefore, they significantly contribute to polyphenol intake, which is commonly considered to be derived only from EPP.

Studies of the metabolic fate of NEPP, although scarce, have shown that NEPP are extensively fermented in the colon, releasing several bioactive and absorbable metabolites.

The main site of action of NEPP is the colon, with several studies of different animal models showing different mechanisms by which NEPP may play a preventive role with regard to colorectal cancer. Also, NEPP (through their bioactive metabolites) may also play a role in the prevention

of other chronic diseases, in particular in mitigating certain risk factors for CVD.

The health properties of DF and phenolic compounds are commonly attributed to polysaccharides and EPP, respectively; in both cases this disregards the relevant contribution of NEPP to such properties.

Further research into NEPP should be encouraged, including analysis of food content, estimates of dietary intakes in different populations, studies of their metabolic fate and sites of action, and evaluation of their health effects through clinical trials. This would help us to gain a complete overview of the biological and nutritional relevance of this understudied fraction of dietary antioxidants.

### Acknowledgements

J. P.-J. acknowledges the Spanish Ministry of Science and Innovation for granting her a 'Juan de la Cierva' postdoctoral contract. M. E. D.-R. acknowledges the CSIC (Spanish Research Council) for granting her a 'JAE-Doc' postdoctoral contract. The present review was supported by the Spanish Ministry of Science and Innovation (project AGL2011–27741). The Spanish Ministry of Science and Innovation and the CSIC had no role in the design, analysis or writing of this article.

F. S.-C. planned the review and designed the manuscript, jointly with J. P.-J. The first version of the manuscript was written by J. P.-J., supervised by F. S.-C. All the authors contributed to writing the manuscript and approved the final version.

None of the authors had any conflict of interest.

### References

- Brighenti F, Valtueña S, Pellegrini N, *et al.* (2005) Total antioxidant capacity of the diet is inversely and independently related to plasma concentration of high-sensitivity C-reactive protein in adult Italian subjects. *Br J Nutr* **93**, 619–625.
- Valtueña S, Pellegrini N, Franzini L, *et al.* (2008) Food selection based on total antioxidant capacity can modify antioxidant intake, systemic inflammation, and liver function without altering markers of oxidative stress. *Am J Clin Nutr* **87**, 1290–1297.
- Del Rio D, Agnoli C, Pellegrini N, *et al.* (2011) Total antioxidant capacity of the diet is associated with lower risk of ischemic stroke in a large Italian cohort. *J Nutr* **141**, 118–123.
- Franzini L, Ardigo D, Valtueña S, *et al.* (2012) Food selection based on high total antioxidant capacity improves endothelial function in a low cardiovascular risk population. *Nutr Metab Cardiovasc Dis* **22**, 50–57.
- Serafini M, Jakszyn P, Luján-Barroso L, *et al.* (2012) Dietary total antioxidant capacity and gastric cancer risk in the European Prospective Investigation into Cancer and Nutrition study. *Int J Cancer* **131**, E544–E554.
- Bravo L, Abia R & Saura-Calixto F (1994) Polyphenols as dietary fibre associated compounds. Comparative study on *in vivo* and *in vitro* properties. *J Agric Food Chem* **42**, 1481–1487.

7. Sofi F, Cesari F, Abbate R, *et al.* (2008) Adherence to Mediterranean diet and health status: meta-analysis. *BMJ* **337**, 673–675.
8. Halliwell B (1996) Antioxidants in human health and disease. *Ann Rev Nutr* **16**, 33–50.
9. O'Neill ME, Carroll Y, Corridan B, *et al.* (2001) A European carotenoid database to assess carotenoid intakes and its use in a five-country comparative study. *Br J Nutr* **85**, 499–507.
10. Saura-Calixto F & Goñi I (2006) Antioxidant capacity of the Spanish Mediterranean diet. *Food Chem* **94**, 442–447.
11. Manach C, Williamson G, Morand C, *et al.* (2005) Bioavailability and bioefficacy of polyphenols in humans. I. Review of 97 bioavailability studies. *Am J Clin Nutr* **81**, 230S–242S.
12. Ramos S (2007) Effects of dietary flavonoids on apoptotic pathways related to cancer chemoprevention. *J Nutr Biochem* **18**, 427–442.
13. Scalbert A, Manach C, Morand C, *et al.* (2005) Dietary polyphenols and the prevention of diseases. *Crit Rev Food Sci Nutr* **45**, 287–306.
14. Hooper L, Kroon PA, Rimm EB, *et al.* (2008) Flavonoids, flavonoid-rich foods, and cardiovascular risk: a meta-analysis of randomized controlled trials. *Am J Clin Nutr* **88**, 38–50.
15. Rossi M, Bosetti C, Negri E, *et al.* (2010) Flavonoids, proanthocyanidins, and cancer risk: a network of case-control studies from Italy. *Nutr Cancer* **62**, 871–877.
16. Pérez-Jiménez J & Torres JL (2011) Analysis of non-extractable phenolic compounds in foods: the current state of the art. *J Agric Food Chem* **59**, 12713–12724.
17. Le Bourvellec C, Bouchet B & Renard CMGC (2005) Non-covalent interaction between procyanidins and apple cell wall material. Part III: Study on model polysaccharides. *Biochim Biophys Acta* **1725**, 10–18.
18. Faulds CB & Williamson G (1999) The role of hydroxycinnamates in the plant cell wall. *J Sci Food Agric* **79**, 393–395.
19. Burr SJ & Fry SC (2009) Extracellular cross-linking of maize arabinoxylans by oxidation of feruloyl esters to form oligoferuloyl esters and ether-like bonds. *Plant J* **58**, 554–567.
20. Bunzel M (2010) Chemistry and occurrence of hydroxycinnamate oligomers. *Phytochem Rev* **9**, 47–64.
21. Matthews S, Mila I, Scalbert A, *et al.* (1997) Extractable and non-extractable proanthocyanidins in barks. *Phytochemistry* **45**, 405–410.
22. Saura-Calixto F (1987) Dietary fibre complex in a sample rich in condensed tannins and uronic acids. *Food Chem* **23**, 95–103.
23. Saura-Calixto F, Goni I, Manas E, *et al.* (1991) Klason lignin, condensed tannins and resistant protein as dietary fibre constituents: determination in grape pomaces. *Food Chem* **39**, 299–309.
24. Vitaglione P, Napolitano A & Fogliano V (2008) Cereal dietary fibre: a natural functional ingredient to deliver phenolic compounds into the gut. *Trends Food Sci Technol* **19**, 451–463.
25. Palafox-Carlos H, Ayala-Zavala JF & González-Aguilar GA (2011) The role of dietary fiber in the bioaccessibility and bioavailability of fruit and vegetable antioxidants. *J Food Sci* **76**, R6–R15.
26. Saura-Calixto F (2011) Dietary fibre as a carrier of dietary antioxidants: an essential physiological function. *J Agric Food Chem* **59**, 43–49.
27. Pérez-Jiménez J, Arranz S & Saura-Calixto F (2009) Proanthocyanidin content in foods is largely underestimated in the literature data: an approach to quantification of the missing proanthocyanidins. *Food Res Intl* **42**, 1381–1388.
28. Saura-Calixto F, Serrano J & Goñi I (2007) Intake and bioaccessibility of total polyphenols in a whole diet. *Food Chem* **101**, 492–501.
29. Arranz S, Saura-Calixto F, Shaha S, *et al.* (2009) High contents of nonextractable polyphenols in fruits suggest that polyphenol contents of plant foods have been underestimated. *J Agric Food Chem* **57**, 7298–7303.
30. Rufino MdSM, Pérez-Jiménez J, Arranz S, *et al.* (2011) Açai (*Euterpe oleraceae*) 'BRS Pará': a tropical fruit source of antioxidant dietary fiber and high antioxidant capacity oil. *Food Res Intl* **44**, 2100–2106.
31. Guyot S, Le Bourvellec C, Marnet N, *et al.* (2002) Procyanidins are the most abundant polyphenols in dessert apples at maturity. *LWT-Food Sci Technol* **35**, 289–291.
32. Hamauzu Y & Mizuno Y (2011) Non-extractable procyanidins and lignin are important factors in the bile acid binding and radical scavenging properties of cell wall material in some fruits. *Plant Foods Human Nutr* **66**, 70–77.
33. Tow WW, Premier R, Jing H, *et al.* (2011) Antioxidant and antiproliferation effects of extractable and nonextractable polyphenols isolated from apple waste using different extraction methods. *J Food Sci* **76**, 163–172.
34. White BL, Howard LR & Prior RL (2010) Release of bound procyanidins from cranberry pomace by alkaline hydrolysis. *J Agric Food Chem* **58**, 7572–7579.
35. Hellström JK, Törrönen AR & Mattila PH (2009) Proanthocyanidins in common food products of plant origin. *J Agric Food Chem* **57**, 7899–7906.
36. Saura Calixto F (1988) Effect of condensed tannins in the analysis of dietary fiber in carob pods. *J Food Sci* **53**, 1769–1771.
37. Goñi I, Díaz-Rubio ME, Pérez-Jiménez J, *et al.* (2009) Towards an updated methodology for measurement of dietary fiber, including associated polyphenols, in food and beverages. *Food Res Intl* **42**, 840–846.
38. Bianco A & Uccella N (2000) Biophenolic components of olives. *Food Res Intl* **33**, 475–485.
39. Gruz J, Ayaz FA, Torun H, *et al.* (2011) Phenolic acid content and radical scavenging activity of extracts from medlar (*Mespilus germanica* L.) fruit at different stages of ripening. *Food Chem* **124**, 271–277.
40. Ye XQ, Chen JC, Liu DH, *et al.* (2011) Identification of bioactive composition and antioxidant activity in young mandarin fruits. *Food Chem* **124**, 1561–1566.
41. Rufino MdSM, Pérez-Jiménez J, Tabernero M, *et al.* (2010) Acerola and cashew apple as sources of antioxidants and dietary fibre. *Int J Food Sci Technol* **45**, 2227–2233.
42. Kapasakalidis PG, Rastall RA & Gordon MH (2006) Extraction of polyphenols from processed black currant (*Ribes nigrum* L.) residues. *J Agric Food Chem* **54**, 4016–4021.
43. Jung MY, Jeon BS & Bock JY (2002) Free, esterified, and insoluble-bound phenolic acids in white and red Korean ginsengs (*Panax ginseng* C.A. Meyer). *Food Chem* **79**, 105–111.
44. Krygier K, Sosulski F & Hogge L (1982) Free, esterified and insoluble-bound phenolic acids. 2. Composition of phenolic acids in rapeseed flour and hulls. *J Agric Food Chem* **30**, 334–336.
45. Adom KK & Liu RH (2002) Antioxidant activity of grains. *J Agric Food Chem* **50**, 6182–6187.
46. Hatcher DW & Kruger JE (1997) Simple phenolic acids in flours prepared from Canadian wheat: relationship to ash content, color, and polyphenol oxidase activity. *Cereal Chem* **74**, 337–343.
47. Gallardo C, Jiménez L & García-Conesa MT (2006) Hydroxycinnamic acid composition and *in vitro* antioxidant activity of selected grain fractions. *Food Chem* **99**, 455–463.
48. Hegde S, Kavitha S, Varadaraj MC, *et al.* (2006) Degradation of cereal bran polysaccharide-phenolic acid complexes by *Aspergillus niger* CFR 1105. *Food Chem* **96**, 14–19.

49. Holtekjolen AK, Kinitz C & Knutsen SH (2006) Flavanol and bound phenolic acid contents in different barley varieties. *J Agric Food Chem* **54**, 2253–2260.
50. Hosseini FS & Mazza G (2009) Triticale bran and straw: potential new sources of phenolic acids, proanthocyanidins, and lignans. *J Functional Foods* **1**, 57–64.
51. Chandrasekara A & Shahidi F (2010) Content of insoluble bound phenolics in millets and their contribution to antioxidant capacity. *J Agric Food Chem* **58**, 6706–6714.
52. John JA & Shahidi F (2010) Phenolic compounds and antioxidant activity of Brazil nut (*Bertholletia excelsa*). *J Functional Foods* **2**, 196–209.
53. Li L, Tsao R, Yang R, *et al.* (2006) Polyphenolic profiles and antioxidant activities of heartnut (*Juglans ailanthifolia* var. *cordiformis*) and Persian walnut (*Juglans regia* L.). *J Agric Food Chem* **54**, 8033–8040.
54. Zurita J, Díaz-Rubio ME & Saura-Calixto F (2012) Improved procedure to determine non-extractable polymeric proanthocyanidins in plant foods. *Intl J Food Sci Nutr* **63**, 936–939.
55. United States Department of Agriculture (2004) USDA Database for the proanthocyanidin content of selected foods. <http://www.ars.usda.gov/nutrientdata> (accessed September 2012).
56. Arranz S, Silván JM & Saura-Calixto F (2010) Nonextractable polyphenols, usually ignored, are the major part of dietary polyphenols: a study on the Spanish diet. *Molec Nutr Food Res* **54**, 1646–1658.
57. Arranz S, Pérez-Jiménez J & Saura-Calixto F (2008) Antioxidant capacity of walnut (*Juglans regia* L.): contribution of oil and defatted matter. *Eur Food Res Technol* **227**, 425–431.
58. Hervet-Hernández D, García OP, Rosado JL, *et al.* (2011) The contribution of fruits and vegetables to dietary intake of polyphenols and antioxidant capacity in a Mexican rural diet: importance of fruit and vegetable variety. *Food Res Intl* **44**, 1182–1189.
59. Kroon PA, Faulds CB, Ryden P, *et al.* (1997) Release of covalently bound ferulic acid from fiber in the human colon. *J Agric Food Chem* **45**, 661–667.
60. Rondini L, Peyrat-Maillard MN, Marsset-Baglieri A, *et al.* (2004) Bound ferulic acid from bran is more bioavailable than the free compound in rat. *J Agric Food Chem* **52**, 4338–4343.
61. Pérez-Jiménez J, Serrano J, Tabernero M, *et al.* (2009) Bioavailability of phenolic antioxidants associated with dietary fiber: plasma antioxidant capacity after acute and long-term intake in humans. *Plant Foods Hum Nutr* **64**, 102–107.
62. Touriño S, Fuguet E, Vinardell MP, *et al.* (2009) Phenolic metabolites of grape antioxidant dietary fiber in rat urine. *J Agric Food Chem* **57**, 11418–11426.
63. Saura-Calixto F, Pérez-Jiménez J, Touriño S, *et al.* (2010) Proanthocyanidin metabolites associated with dietary fibre from *in vitro* colonic fermentation and proanthocyanidin metabolites in human plasma. *Molec Nutr Food Res* **54**, 939–946.
64. Touriño S, Pérez-Jiménez J, Mateos-Martín ML, *et al.* (2011) Metabolites in contact with the rat digestive tract after ingestion of a phenolic-rich dietary fiber matrix. *J Agric Food Chem* **59**, 5955–5963.
65. Mateos-Martín ML, Pérez-Jiménez J, Fuguet E, *et al.* (2012) Non-extractable proanthocyanidins from grape are a source of bioavailable (epi)catechin and derived metabolites in rats. *Br J Nutr* **108**, 290–297.
66. Braune A, Bunzel M, Yonekura R, *et al.* (2009) Conversion of dehydrodiferulic acids by human intestinal microbiota. *J Agric Food Chem* **57**, 3556–3562.
67. Cerdá B, Espín JC, Parra S, *et al.* (2004) The potent *in vitro* antioxidant ellagitannins from pomegranate juice are metabolised into bioavailable but poor antioxidant hydroxy-6H-dibenzopyran-6-one derivatives by the colonic microflora of healthy humans. *Eur J Nutr* **43**, 205–220.
68. Crozier A, Jaganath IB & Clifford MN (2009) Dietary phenolics: chemistry, bioavailability and effects on health. *Nat Prod Rep* **26**, 1001–1043.
69. Selma MV, Espín JC & Tomás-Barberán FA (2009) Interactions between phenolics and gut microbiota: role in human health. *J Agric Food Chem* **57**, 6485–6501.
70. Aprikian O, Duclos V, Guyot S, *et al.* (2003) Apple pectin and a polyphenol-rich apple concentrate are more effective together than separately on cecal fermentations and plasma lipids in rats. *J Nutr* **133**, 1860–1865.
71. Juśkiewicz J, Milala J, Jurgonński A, *et al.* (2011) Consumption of polyphenol concentrate with dietary fructo-oligosaccharides enhances cecal metabolism of quercetin glycosides in rats. *Nutrition* **27**, 351–357.
72. Schroeter H, Heiss C, Balzer J, *et al.* (2006) (–)-Epicatechin mediates beneficial effects of flavanol-rich cocoa on vascular function in humans. *PNAS* **103**, 1024–1029.
73. Karlsson PC, Huss U, Jenner A, *et al.* (2005) Human fecal water inhibits COX-2 in colonic HT-29 cells: role of phenolic compounds. *J Nutr* **135**, 2343–2349.
74. Larrosa M, Luceri C, Vivoli E, *et al.* (2009) Polyphenol metabolites from colonic microbiota exert anti-inflammatory activity on different inflammation models. *Mol Nutr Food Res* **53**, 1044–1054.
75. Verzelloni E, Pellacani C, Tagliazucchi D, *et al.* (2011) Antigliative and neuroprotective activity of colon-derived polyphenol catabolites. *Molec Nutr Food Res* **55**, S35–S43.
76. González-Sarriás A, Espín JC, Tomás-Barberán FA, *et al.* (2009) Gene expression, cell cycle arrest and MAPK signalling regulation in Caco-2 cells exposed to ellagic acid and its metabolites, urolithins. *Molec Nutr Food Res* **53**, 686–698.
77. Velázquez OC, Lederer HM & Rombeau JL (1996) Butyrate and the colonocyte: implications for neoplasia. *Dig Dis Sci* **41**, 727–739.
78. Wong JMW, De Souza R, Kendall CWC, *et al.* (2006) Colonic health: fermentation and short chain fatty acids. *J Clin Gastroenterol* **40**, 235–243.
79. Hosseini E, Grootaert C, Verstraete W, *et al.* (2011) Propionate as a health-promoting microbial metabolite in the human gut. *Nutr Rev* **69**, 245–258.
80. Bravo L, Saura-Calixto F & Goni I (1992) Effects of dietary fibre and tannins from apple pulp on the composition of faeces in rats. *Br J Nutr* **67**, 463–473.
81. Martín-Carrón N, García-Alonso A, Goñi I, *et al.* (1997) Nutritional and physiological properties of grape pomace as a potential food ingredient. *Am J Enol Vitic* **48**, 328–332.
82. Martín-Carrón N, Saura-Calixto F & Goñi I (2000) Effects of dietary fibre- and polyphenol-rich grape products on lipidaemia and nutritional parameters in rats. *J Sci Food Agric* **80**, 1183–1188.
83. Bravo L, Mañas E & Saura-Calixto F (1993) Dietary non-extractable condensed tannins as indigestible compounds: effects on faecal weight and protein and fat excretion. *J Sci Food Agric* **63**, 63–68.
84. Goñi I, Jiménez-Escrig A, Gudiel M, *et al.* (2005) Artichoke (*Cynara scolymus* L) modifies bacterial enzymatic activities and antioxidant status in rat cecum. *Nutr Res* **25**, 607–615.
85. Goñi I & Serrano J (2005) The intake of dietary fiber from grape seeds modifies the antioxidant status in rat cecum. *J Sci Food Agric* **85**, 1877–1881.
86. López-Oliva ME, Pozuelo MJ, Rotger R, *et al.* (2013) Grape antioxidant dietary fibre prevents mitochondrial apoptotic pathways by enhancing Bcl-2 and Bcl-xL expression and minimising oxidative stress in rat distal colonic mucosa. *Br J Nutr* **109**, 4–16.



87. Lizarraga D, Vinardell MP, Noé V, *et al.* (2011) A lyophilized red grape pomace containing proanthocyanidin-rich dietary fiber induces genetic and metabolic alterations in colon mucosa of female C57Bl/6J mice. *J Nutr* **141**, 1597–1604.
88. Sánchez-Tena S, Lizarraga D, Miranda A, *et al.* (2013) Grape antioxidant dietary fiber inhibits intestinal polyposis in ApcMin/+ mice: relation to cell cycle and immune response. *Carcinogenesis* (publication ahead of print version 29 May 2013).
89. López-Oliva ME, Agis-Torres A, García-Palencia P, *et al.* (2006) Induction of epithelial hypoplasia in rat caecal and distal colonic mucosa by grape antioxidant dietary fiber. *Nutr Res* **26**, 651–658.
90. López-Oliva ME, Agis-Torres A, Goñi I, *et al.* (2010) Grape antioxidant dietary fibre reduced apoptosis and induced a pro-reducing shift in the glutathione redox state of the rat proximal colonic mucosa. *Br J Nutr* **103**, 1110–1117.
91. Pozuelo MJ, Agis-Torres A, Hervert-Hernández D, *et al.* (2012) Grape antioxidant dietary fibre stimulates *Lactobacillus* growth in rat cecum. *J Food Sci* **77**, H59–H62.
92. Hervert-Hernández D, Pintado C, Rotger R, *et al.* (2009) Stimulatory role of grape pomace polyphenols on *Lactobacillus acidophilus* growth. *Int J Food Microbiol* **136**, 119–122.
93. Pérez-Jiménez J, Serrano J, Tabernero M, *et al.* (2008) Effects of grape antioxidant dietary fiber in cardiovascular disease risk factors. *Nutrition* **24**, 646–653.
94. Vitaglione P & Fogliano V (2010) Cereal fibres, antioxidant activity and health. In *Dietary Fibre: New Frontiers for Food and Health*, pp. 379–390 [JW Van Der Kamp, J Jones, B McCleary and D Topping, editors]. Wageningen: Wageningen Academic Publishers.
95. Robertson MD (2007) Metabolic cross talk between the colon and the periphery: implications for insulin sensitivity. Sir David Cuthbertson Medal Lecture. *Proc Nutr Soc* **66**, 351–361.
96. Lecumberri E, Goya L, Mateos R, *et al.* (2007) A diet rich in dietary fiber from cocoa improves lipid profile and reduces malondialdehyde in hypercholesterolemic rats. *Nutrition* **23**, 332–341.
97. Cracowski JL & Ormezzano O (2004) Isoprostanes, emerging biomarkers and potential mediators in cardiovascular diseases. *Eur Heart J* **25**, 1675–1678.